

Regional climate downscaling with prior statistical correction of the global climate forcing

Augustin Colette, Robert Vautard, M. Vrac

▶ To cite this version:

Augustin Colette, Robert Vautard, M. Vrac. Regional climate downscaling with prior statistical correction of the global climate forcing. Geophysical Research Letters, 2012, 39 (13), pp.NC. 10.1029/2012GL052258. ineris-00961787

HAL Id: ineris-00961787 https://ineris.hal.science/ineris-00961787

Submitted on 20 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Regional climate downscaling with prior statistical correction of the global climate

2 forcing

3 A. Colette (1), R. Vautard (2), M. Vrac (2)

Institut National de l'Environnement Industriel et des Risques, INERIS, Verneuil-en Halatte, France

6 2. Laboratoire des Sciences du Climat et de l'Environnement / Institut Pierre-Simon Laplace

7 (LSCE/IPSL), laboratoire CEA/CNRS/UVSQ, Gif-sur-Yvette, France

8 Corresponding author address : augustin.colette@ineris.fr

9 Abstract

10 A novel climate downscaling methodology that attempts to correct climate simulation biases is proposed. By combining an advanced statistical bias correction method with a dynamical 11 downscaling it constitutes a hybrid technique that yields nearly unbiased, high-resolution, 12 13 physically consistent, three-dimensional fields that can be used for climate impact studies. The method is based on a prior statistical distribution correction of large-scale global climate 14 15 model (GCM) 3-dimensional output fields to be taken as boundary forcing of a dynamical 16 regional climate model (RCM). GCM fields are corrected using meteorological reanalyses. We evaluate this methodology over a decadal experiment. The improvement in terms of 17 18 spatial and temporal variability is discussed against observations for a past period. The biases 19 of the downscaled fields are much lower using this hybrid technique, up to a factor 4 for the mean temperature bias compared to the dynamical downscaling alone without prior bias 20 correction. Precipitation biases are subsequently improved hence offering optimistic 21 22 perspectives for climate impact studies.

23 **1 Introduction**

Global Climate Models (GCM) improved notably in their representation of the climate system
over the past couple of decades (IPCC, 2007). Their design is focused on the global scale, and

their main scope consists in capturing the sensitivity of the global climate to changes in external natural and anthropogenic forcing. The fairly low resolution of such models does not allow for the detailed simulation of local atmospheric processes. In addition, the main focus being the global energy balance, coupled models may exhibit significant regional biases in important variables such as temperature or precipitation.

However, climate risk assessment requires horizontal resolution of the order of half a degree or below and unbiased projections, especially when it comes to meteorological extremes. More generally such information is required in order to design adaptation measures for which impact models (e.g., with regards to food safety, energy, water, air pollution), tuned on current climate observations, need to be applied to future climate projections. Such a requirement cannot be met by current raw GCM outputs.

The transformation of global model outputs into high spatial resolution products is referred to 37 38 as climate downscaling. It can be divided into two broad types of approaches: statistical or dynamical downscaling. Statistical downscaling builds upon a prior knowledge of statistical 39 relationships between the GCM and monitoring data. Statistical models representing those 40 relationships are then applied over future time periods, without involving any additional 41 physical modelling in addition to the GCM (Wilks and Wilby, 1999, Vrac et al., 2007, 42 Semenov et al., 1998, Maraun et al., 2010). To downscale a global model in a dynamical way, 43 one implements a Regional Climate Model (RCM) forced by the global fields at the 44 boundaries (Giorgi et al., 2009, Laprise, 2008). Similarly to the GCM, the RCM provides a 45 comprehensive physically-consistent representation of the climate system. However, GCM 46 biases are conveyed to the RCM, and the latter can only compensate, or enhance, these flaws. 47 In order to cope with these deficiencies, bias correction methods are often applied to RCM 48 outputs prior to the implementation of an impact model (Christensen et al., 2008, Oettli et al., 49 2011). However this methodology suffers from several caveats. On the one hand, the fields 50

are generally corrected without considering spatial, temporal or inter-variable correlation. On 51 52 the other hand, the bias correction requires high-resolution observations, generally not available on a grid, but rather at scattered locations. These problems could be at least partly 53 avoided if most of the GCM biases were removed before the dynamical downscaling, an 54 approach that we investigate in this article. A few studies investigated the possibility to 55 correct large scale forcing prior to applying a mesoscale model (Rasmussen et al., 2012, Schär 56 et al., 1996) but none of them achieved that with a downscaling technique that matches the 57 whole range of the distribution to meteorological reanalyses. 58

We propose here an innovative downscaling methodology that combines both dynamical and statistical approaches, but in a different order compared to what is usually done. In a nutshell, our hybrid approach consists in applying a statistical correction of the GCM fields with respect to atmospheric reanalyses prior to performing a dynamical downscaling of these corrected fields. As such, this approach constitutes a hybrid climate downscaling technique building upon upstream statistical correction and downstream physical modelling.

Like any probabilistic downscaling technique, the upstream statistical correction may alter the 65 integrity of the forcing fields by matching it to reanalyses. The main strength of our hybrid 66 approach lies in the implementation of a mesoscale model after the probabilistic downscaling 67 that guarantees the physical consistency of the resulting fields and hence constitutes an 68 essential advantage for climate impact studies (Parry et al., 2007). Statistical downscaling that 69 targets only a couple of surface variables has long been considered satisfactory for most 70 climate impact studies (such as food safety or hydrological extremes). However other 71 72 applications such as air quality modelling require physically-consistent 3D atmospheric fields. That is why regional air quality projection studies rely on raw RCM outputs, and our 73 74 technique offers a unique perspective to derive unbiased, balanced, 3D forcing fields.

In section 2, the statistical and physical downscaling methodologies are presented. The evaluation results are given in section 3 on a test case for present day simulation. The application to future projections is left for upcoming studies.

78 2 Methodology

79 2.1 Large scale climate model

The large scale climate model that we use to demonstrate the efficiency of our hybrid statistical and dynamical technique is the coupled climate model IPSLcm (Institut Pierre Simon Laplace Coupled Model) GCM (Marti et al., 2010).

The simulations used here are obtained with the "low resolution" versions prepared for the CMIP5 (Climate Model Intercomparison Project) stream of the Intergovernmental Panel on Climate Change (IPCC). The meteorological fields are computed on a global 96x96 points grid with a horizontal resolution of 3.75 x 1.875 degrees and 39 vertical levels.

87 2.2 Statistical downscaling

The probabilistic downscaling methodology used here is the CDF-t (Cumulative Distribution 88 Function transform) of (Michelangeli et al., 2009), based on a variant of the "quantile-89 matching" technique (Déqué, 2007). Quantile-matching consists in associating to a modelled 90 value, the value in a control distribution (e.g. observations) that has the same probability. In 91 other words, from a quantile in the CDF of the simulations, the corresponding quantile in the 92 CDF of the control data (e.g. observations) is determined. By scaling the quantile-quantile 93 relationship, the correction changes the shape of the distribution so that the events whose 94 frequency (or probability) is systematically biased in the model are better captured. 95

While classical applications of quantile-matching consider that the CDF of the simulations is stationary in time (Maraun et al., 2010, Wilks and Wilby, 1999), the scope of CDF-t consists in expanding this technique for the case where the CDF of the simulations for the future has changed. This is done, first, by estimating the CDF of the corrected variable for the future time period of interest (Michelangeli et al., 2009). Then, projections are obtained through a quantile-quantile technique between future uncorrected and corrected CDFs (Vrac et al., 2012). The methodology implemented here thus applies for future projections even though we decided to limit the scope of the present paper to historical periods in order to discuss its validation.

105 This CDF-t technique has been used successfully in the past to downscale climate models (Vrac et al., 2012, Flaounas et al., 2011, Michelangeli et al., 2009) but one should note the 106 107 two major limitations of the approach. First, only the bulk CDF is matched, the temporal frequency and spatial patterns are not altered so that any flaw in the persistence or in the 108 109 spatial distribution of the weather patterns is not improved. In addition, the major underlying hypothesis of the CDF-t downscaling is that, although the CDFs are not supposed to be 110 stationary, the transformation T from model to observed variable CDFs is supposed to be 111 112 valid under changed climate conditions, i.e. is supposed stationary in time. We emphasize that even though CDF-t is designed to be applied to future climate simulations, we decided to 113 114 apply this technique in the present paper to a current period for validation purposes.

115 2.3 Dynamical downscaling

We use the Weather Research and Forecasting (Skamarock et al., 2008) mesoscale model to downscale the IPSLcm fields in a dynamical way. The spatial resolution is 50km and the domain covers the whole of Europe with 119x116 grid points. The setup is the same as that of (Menut et al., 2012) who present a detailed evaluation of the performance of the IPSLcm/WRF regional climate modelling suite. However no nudging was applied in the present case in order to evaluate the full effect of prior correction on dynamical downscaling.

122 2.4 Experimental design

We perform a CDF-t based correction of the large-scale input fields produced with theIPSLcm model so that corrected fields will be used for the dynamical downscaling.

Distributions are matched with those of reanalysed fields of the ERA-interim reanalysis. Unlike existing applications of CDF-t that perform a scaling of large-scale model outputs to point surface observations (Michelangeli et al., 2009) or gridded surface analyses (Flaounas et al., 2011) we scale several variables of the model to the whole 3D fields of the reanalysis.

The correction is achieved at each GCM grid-point independently, where reanalysed fields 129 were previously interpolated. There was no attempt to maintain the spatial consistency of the 130 fields considering that (1) matched fields are coarse enough to avoid the introduction of high-131 frequency variability and (2) potential spurious features would vanish after having used the 132 mesoscale model to downscale the corrected fields. For each variable and at each grid point, 133 134 we extract the time series for the whole period to produce the two distributions (GCM and reanalysis) that will be matched. To account for seasonality, all training distributions are taken 135 on a monthly basis. For 3D and surface temperature, the correction is performed 136 independently for the 4 daily time steps to account for the diurnal cycle. Since we match the 137 bulk distribution of the time series, there is no matching of sequences of event, on the 138 contrary we maintain the temporal consistency of the input field. 139

140 The correction is done for 3D zonal and meridional wind, 3D relative humidity, and 3D and surface (skin) temperature. Surface pressure and geopotential height are not matched in order 141 to maintain flow consistency and quasi-geostrophy at the boundaries, but they are indirectly 142 modified by the matching of the 3D temperature field. The hydrostatic balance of the 143 corrected input field is recomputed before launching the mesoscale model in order to ensure 144 physical consistency along the columns; by proceeding to an upward integration of the 145 hydrostatic balance, corrections applied to the temperature field are conveyed to the 146 geopotential height. 147

The evaluation experiment consists of simulations over a 11-year period for the downscaling.The first year is considered as a spin-up period and it is thus discarded from the following

analysis. The last decade of the 20th century is chosen because of the full overlap between 150 ERA-interim and IPCC historical simulations. This time period also allows comparing the 151 efficiency of the methodology against observations. Two simulations are carried out, starting 152 on 1 January 1989. The first one is done without applying the GCM correction prior to 153 dynamical downscaling, while the second is done with application of the prior CDF-t 154 approach. The two simulations are then compared to E-OBS data (Haylock et al., 2008) over 155 the same time period. Since the focus of this study is not to validate the performance of the 156 157 CFD-t itself directly applied to the GCM fields (as it was demonstrated before (Flaounas et al., 2011, Michelangeli et al., 2009, Vrac et al., 2012)), but the impact of CDF-t on the 158 dynamical regional climate downscaling, it was unnecessary to implement a 'leave-one-out' 159 testing approach. The duration of the simulations (10 years) is too short to address the benefits 160 for meteorological extremes; this aspect is left for future work while we focus here on average 161 162 biases.

163 **3 Results**

164 The evaluation of the results is performed against the European Climate Gridded dataset (E-165 OBS) temperature and precipitation observations.

166 **3.1 Surface temperature**

167 The bias of temperature averaged over the 10-year time period is given in Figure 1 for the reanalysis (ERA-i), the large-scale climate model (IPSLcm) and its statistically corrected 168 version, the dynamically downscaled climate model (IPSLcm/WRF) and the hybrid 169 statistical/dynamical downscaling (IPSLcm/CDF-t/WRF). For all the models the temperature 170 is interpolated at 950hPa while the observations are provided at 2-m altitude. The 171 discrepancies between E-OBS and ERA-i are confined to the outskirts of the domain where 172 the gap filling procedure used in E-OBS has uncertainties as a result of the scarcity of the 173 monitoring network. In addition, important differences are found over mountainous areas due 174

to lack of resolution and methodological differences. On average, the difference between 175 ERA-i and the observations is -1.41K (standard deviation σ =2.03) over the Western part of 176 the domain (5W, 15E, 40N, 55N). Raw GCM temperatures exhibit a strong negative bias 177 (-4.78K, σ =0.6), except over mountainous areas where the positive biases result from an 178 179 artefact of the smooth orography. This strong negative bias of the low resolution version of the IPSLcm model was discussed before (Hourdin et al., 2012) and was improved in a more 180 recent version of the model including a higher resolution (Cattiaux et al., 2012). This feature 181 constitutes a somewhat good test case for the hybrid downscaling methodology presented 182 here. The statistical correction is efficient at reducing the temperature bias of IPSLcm, the 183 average bias of the corrected GCM is -1.36 (σ =2.07) and its pattern resembles that of ERA-i. 184 The negative bias of IPSLcm is amplified in the raw regional climate model simulations 185 186 (-5.06K, σ =1.49), as was observed by (Menut et al., 2012). The dynamical downscaling does not constrain the distribution in any ways, and it appears that a negative feedback occurs 187 here as the RCM increases the negative biases of forcing fields. On the contrary, the situation 188 189 is better for the hybrid downscaling, the average bias is limited to -2.33K (σ =1.35). The mesoscale still tends to cool down the GCM, and the average bias is larger than for the 190 191 corrected version of IPSLcm since the compensation that occurred over high elevation terrain vanishes. Despite the reduction of the mean bias, it still exhibits a regional pattern with 192 negative values in Western and Northern areas and positive values in Mediterranean areas. 193

194 The overall negative bias is primarily found for low temperatures during winter and to a lesser 195 extent for warm temperatures, even though a bias remains over the lowermost part of the 196 distribution.

197 Seasonality has a strong impact, the mesoscale model tends to be warmer than the large scale 198 forcing in winter (0.5 and 0.6K average bias for IPSLcm and IPSLcm/CDF-t, respectively) 199 and colder in summer (-0.3 and -1.67K average bias for IPSLcm and IPSLcm/CDF-t, respectively). The upstream statistical correction influences indirectly the atmospheric flow.
This feature is confirmed with average sea-level pressure maps (not shown) that exhibit larger
differences in winter than in summer, explaining this uneven influence on temperature of the
bias correction over the year.

204 3.2 Precipitation

Beyond its relevance for climate impact studies, precipitation is an interesting variable to evaluate our methodology since, unlike temperature, this variable was not directly corrected by the prior statistical CDF-t method. The absolute differences between modelled and observed precipitations are provided on **Figure 2**.

The GCM exhibits an overestimation of precipitations throughout the domain. Only Westfacing coastal areas have a deficit, presumably because of the too coarse resolution that is not able to capture the precipitation local maxima over the coastlines. The overestimation is less pronounced over mountainous areas because of a compensation of errors.

The dynamical downscaling of the raw GCM outputs yields an even stronger overestimation of the precipitation because of a negative feedback related to the low temperature bias. The deficit over coastlines and mountains is compensated by the higher resolution of the model.

It is only with the hybrid downscaling that the results are significantly improved. The model still exhibits an overestimation of precipitation but, over low-lying area of Western Europe, the bias is decreased by a factor of two. An excess is found over the Alps. Precipitation deficits are found around the Mediterranean, the spatial patterns of these deficits do not appear highly correlated to coastlines. It may thus be attributable to other uncorrected deficiencies such as weather regime frequencies rather than resolution issues.

The distribution of daily precipitation shows that the hybrid downscaling constitutes an improvement over the whole range of the distribution. Nevertheless, all the simulations still exhibit an overestimation of low precipitations and an underestimation of higher quantiles.

9

225 **4** Conclusion

226 We propose an innovative climate downscaling methodology that combines state-of-the-art statistical and dynamical approaches. We apply a statistical correction to large-scale fields of 227 a Global Climate Model (GCM) prior to a regional simulation. The statistical correction 228 makes use of the Cumulative Distribution Function transformation (CDF-t) designed by 229 (Michelangeli et al., 2009). The GCM field distributions are matched to those of reanalysed 230 fields in order to apply a correction over the whole 3D domain for several variables. The 231 corrected fields are then provided to a dynamical Regional Climate Model (RCM), so that we 232 can produce bias-corrected, yet physically consistent, 3D fields at higher spatial resolution. 233

An application to present-day climate shows that the statistical upstream correction leads to a reduction of the surface temperature bias of a factor four in the regional climate simulation. This improvement yields, in turn, a lower overestimation of precipitations.

The CDF-t upstream correction does not address yet spatial and temporal variability (climate modes, persistence and weather regimes), the technique remains sensitive to the choice of variables included in the correction and the location of the domain since the forcing is applied at the boundaries. The methodology carries some error compensation mechanisms whose effect is minimised thanks to the implementation of a dynamical downscaling in the lee of the statistical correction.

Nevertheless, considering the magnitude of the improvement in terms of mean bias we conclude that this innovative hybrid statistical/dynamical climate downscaling offers promising perspectives for climate impact studies requiring unbiased, balanced, highresolution 3D fields.

247 Acknowledgements

- 248 The CityZen project, grant agreement 212095 of the European Union's Seventh Framework
- 249 Programme (FP7/2007-2013). The Salut'AIR project of the Primequal Research Programme.
- 250 The CORDEX framework and CEA CCRT computing facilities.

251 **References**

- 252 Cattiaux, J., Douville, H., Ribes, A., Chauvin, F. & Plante, C. (2012): Towards a better
- 253 understanding of changes in wintertime cold extremes over Europe A pilot study with CNRM
- 254 & IPSL atmospheric models. Climate Dynamics under review,
- 255 Christensen, J. H., Boberg, F., Christensen, O. B. & Lucas-Picher, P. (2008): On the need for
- bias correction of regional climate change projections of temperature and precipitation.
- 257 Geophys. Res. Lett. 35/L20709,
- Déqué, M. (2007): Frequency of precipitation and temperature extremes over France in an
 anthropogenic scenario: Model results and statistical correction according to observed values.
 Global and Planetary Change 57/1-2, 16-26.
- Flaounas, E., Drobinski, P., Vrac, M., Bastin, S. & Lebeaupin Brossier, C. (2011): Seasonal
 variability and extremes of precipitation and temperature in the Mediterranean region:
 Evaluation of dynamical and statistical downscaling methods for the frame of HyMeX and
 MED-CORDEX. Climate Dynamics under review,
- Giorgi, F., Jones, C. & Asrar, G. R. (2009): Addressing climate information needs at the
 regional level: the CORDEX framework. WMO Bulletin 58/3, 175-183.
- 267 Haylock, M. R., Hofstra, N., Tank, A. M. G. K., Klok, E. J., Jones, P. D., et al. (2008): A
- European daily high-resolution gridded data set of surface temperature and precipitation for
- 269 1950-2006. Journal of Geophysical Research-Atmospheres 113/D20,
- 270 Hourdin, F., Foujols, M. A., Codron, F., Guemas, V., Dufresne, J., et al. (2012): From
- 271 LMDZ4 to LMDZ5: Impact of the atmospheric model grid configuration on the climate and
- sensitivity of IPSL climate model. Climate Dynamics under review,

- Ipcc (2007): Contribution of Working Group I to the Fourth Assessment Report of theIntergovernmental Panel on Climate Change. In: SOLOMON, S., QIN, D., MANNING, M.,
- 275 CHEN, Z., MARQUIS, M., AVERYT, K. B., TIGNOR, M. & MILLER, H. L. (eds).
- 276 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Laprise, R. (2008): Regional climate modelling. Journal of Computational Physics 227, 3641–
 3666.
- Maraun, D., Wetterhall, F., Ireson, A. M., Chandler, R. E., Kendon, E. J., et al. (2010):
 Precipitation downscaling under climate change. Recent developments to bridge the gap
 between dynamical models and the end user. Reviews of Geophysics 48/3,
- Marti, O., Braconnot, P., Dufresne, J. L., Bellier, J., Benshila, R., et al. (2010): Key features
 of the IPSL ocean atmosphere model and its sensitivity to atmospheric resolution. Climate
 Dynamics 34/1, 1-26.
- Menut, L., Tripathi, O. P., Colette, A., Vautard, R., Flaounas, R., et al. (2012): Evaluation of regional climate model forcing with an air quality perspective. Climate Dynamics under review,
- 288 Michelangeli, P. A., Vrac, M. & Loukos, H. (2009): Probabilistic downscaling approaches:
- Application to wind cumulative distribution functions. Geophys. Res. Lett. 36/11, L11708.
- 290 Oettli, P., Sultan, B., Baron, C. & Vrac, M. (2011): Are regional climate models relevant for
- crop yield prediction in West Africa? Environmental Research Letters 6/1, 014008.
- 292 Parry, M. L., Canziani, O. F., Palutikof, J. P., Van Der Linden, P. J. & Hanson, C. E. (2007):
- 293 Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental
- 294 Panel on Climate Change. In: PRESS, C. U. (eds). Cambridge, UK and New York, USA.
- 295 Rasmussen, R., Liu, C., Ikeda, K., Gochis, D., Yates, D., et al. (2012): High-Resolution
- 296 Coupled Climate Runoff Simulations of Seasonal Snowfall over Colorado: A Process Study
- of Current and Warmer Climate. Journal of Climate 24/12, 3015-3048.

- Schär, C., Frei, C., Lüthi, D. & Davies, H. C. (1996): Surrogate climate change scenarios for
 regional climate models. Geophys. Res. Lett. 23/6, 669-672.
- 300 Semenov, M. A., Brooks, R. J., Barrow, E. M. & Richardson, C. W. (1998): Comparison of
- the WGEN and LARS-WG stochastic weather generators for diverse climates. Climate
 Research 10/2, 95-107.
- 303 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., et al. (2008): A
- 304 Description of the Advanced Research WRF Version 3. In: (eds): Technical Note. NCAR.
- 305 Vrac, M., Drobinski, P., Merlo, A., Herrmann, M., Lavaysse, C., et al. (2012): Dynamical and
- 306 statistical downscaling of the French Mediterranean climate: uncertainty assessment. Natural
- 307 Hazards and Earth System Sciences. under review,
- 308 Vrac, M., Stein, M. & Hayhoe, K. (2007): Statistical downscaling of precipitation through
 309 nonhomogeneous stochastic weather typing. Climate Research 34/3, 169-184.
- 310 Wilks, D. S. & Wilby, R. L. (1999): The weather generation game: a review of stochastic
- weather models. Progress in Physical Geography 23/3, 329-357.
- 312 Figures



Figure 1 : Difference between the mean modelled 950hPa temperature and observed (E-OBS) 2-m temperature (K) over the 1990-1999 decade for ERA-interim, the GCM IPSLcm as well as its corrected version and the RCM WRF driven by raw IPSLcm fields and by downscaled IPSL fields corrected with the CDF-t technique. The green-shaded areas in the WRF field are unavailable because located below the 950hPa level in the hybrid coordinates.

319

313



321

-20

Figure 2 : Same as Figure 1 for the precipitations (mm/day) except that only the results of the



0

10 20 30