



**HAL**  
open science

## New insights into the regulation of *cyp3a65* expression in transgenic *tg(cyp3a65:GFP)* zebrafish embryos

Chedi Erradhouani, Benjamin Piccini, Emmanuelle Maillot-Marechal, Selim Aït-Aïssa, Patrick Balaguer, Xavier Coumoul, François Brion

### ► To cite this version:

Chedi Erradhouani, Benjamin Piccini, Emmanuelle Maillot-Marechal, Selim Aït-Aïssa, Patrick Balaguer, et al.. New insights into the regulation of *cyp3a65* expression in transgenic *tg(cyp3a65:GFP)* zebrafish embryos. *Aquatic Toxicology*, 2025, 279, pp.107250. 10.1016/j.aquatox.2025.107250 . ineris-04930797

**HAL Id: ineris-04930797**

**<https://ineris.hal.science/ineris-04930797v1>**


Submitted on 5 Feb 2025

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## New insights into the regulation of *cyp3a65* expression in transgenic *tg(cyp3a65:GFP)* zebrafish embryos

Chedi Erradhouani<sup>a,b</sup>, Benjamin Piccini<sup>a</sup>, Emmanuelle Maillot-Marechal<sup>a</sup>, Selim Aït-Aïssa<sup>a</sup>, Patrick Balaguer<sup>c</sup>, Xavier Coumoul<sup>b,1,\*</sup>, François Brion<sup>a,\*,1</sup>

<sup>a</sup> Unité écotoxicologie des substances et des milieux, Institut National de l'Environnement Industriel et des Risques (INERIS), 60550 Verneuil-en-Halatte, France

<sup>b</sup> Université Paris Cité, INSERM, Health & Functional Exposomics - HealthFex, 75006 Paris, France

<sup>c</sup> Institut de Recherche en Cancérologie de Montpellier (IRCM), Inserm U1194, Université Montpellier, Institut régional du Cancer de Montpellier (ICM), 34090 Montpellier, France

### ARTICLE INFO

#### Keywords:

Transgenic zebrafish embryos  
CYP3A65  
PXR  
AHR  
GR  
Metabolic endocrine disruptors  
Intestine

### ABSTRACT

Facing the need for alternative models allowing assessment of metabolic-endocrine disrupting chemicals (MDCs), especially in poorly investigated tissues such as the intestine, we recently developed a transgenic zebrafish embryo *in vivo* model, *tg(cyp3a65:GFP)*, expressing the Green Fluorescent Protein (GFP) under the control of the zebrafish *cyp3a65* promoter, ortholog of human *cyp3a4*, a gene coding for a key enzyme of intestinal xenobiotic and endobiotic metabolism. In this study, we aimed to better understand the regulation of *cyp3a65* expression by zfpXR, zfAhR2, and zfGR zebrafish orthologs of well-known human xenosensors PXR and AhR, and steroid nuclear receptor GR. For this purpose, we performed zebrafish embryo *tg(cyp3a65:GFP)* (co)exposures to a variety of agonists (clotrimazole, TCDD, fluticasone propionate) and antagonists (econazole nitrate, CH223181, RU486), which were characterized using *in vitro* zebrafish reporter gene assays. We show that zfpXR and zfAhR2 cooperate to positively regulate *cyp3a65* expression, involving different transcription factors and their interaction. Moreover, for the first time, we show that zfGR agonist strongly inhibits the constitutive expression of *cyp3a65*, and we hypothesized the possible involvement of the transcriptional factor zfHNF4α. These results provide a better understanding of the regulation of zebrafish *cyp3a65* expression, highlighting the complex interaction between different transcription factors, which is consistent with the multiple regulatory pathways of *cyp3a4* in humans. Our data support the idea that this gene is a target of multiple contaminants capable of interacting with zfpXR, zfAhR2 and zfGR and highlights the relevance of the *tg(cyp3a65:GFP)* model to screen chemicals potentially acting as MDCs based on their modes of action at the intestinal level, which could be relevant for hazard assessment of chemicals for human and environmental health.

### Introduction

Among endocrine disrupting chemicals, some referred to metabolic-endocrine disrupting chemicals (MDCs), can promote metabolic changes responsible for an increased prevalence of metabolic diseases in humans such as obesity, diabetes, and/or non-alcoholic fatty liver disease (NAFLD) (Bernal et al., 2022; Petrakis et al., 2017). The development of biological models and bioassays to evaluate metabolic disruptions induced by MDCs has thus become a major challenge in recent years (Kassotis et al., 2022). This challenge meets the need for alternative models to identify modes of action (MoAs) and investigate possible link

with adverse effects (Audouze et al., 2020), especially in poorly investigated organs such as the intestine (Erradhouani et al., 2024).

In this context, we recently established a new *in vivo* transgenic model, *tg(cyp3a65:GFP)* (Erradhouani et al., in submission), developed in pigment-free zebrafish Casper mutant (White et al., 2008). This transgenic line expresses the green fluorescent protein (GFP) under the control of the zebrafish *cyp3a65* gene promoter coding for cytochrome P450 (CYP) CYP3A65, a zebrafish ortholog of human CYP3A4 (Goldstone et al., 2010). The CYP3A4 enzyme plays a central role in intestinal metabolism, as it is involved in the endogenous metabolism such as bile acids, and the protection of the organism against orally

\* Corresponding authors.

E-mail addresses: [xavier.coumoul@u-paris.fr](mailto:xavier.coumoul@u-paris.fr) (X. Coumoul), [francois.brion@ineris.fr](mailto:francois.brion@ineris.fr) (F. Brion).

<sup>1</sup> Equal contribution.

absorbed xenobiotics, catalyzing one of the first steps of their detoxification processes (Chen et al., 2014; van Herwaarden et al., 2009; van Waterschoot et al., 2009). Previous results showed that *cyp3a65* expression is regulated by two xenobiotic receptors, zfpXR and zfAhR2, orthologs of human PXR (pregnane X receptor) and AhR (aryl hydrocarbon receptor), two well-known ligand-activated transcription factors (Chang et al., 2013; Kubota et al., 2015). Moreover, interactions or crosstalk between zfpXR and zfAhR2 in the regulation of *cyp3a65* have been suggested but are not fully understood. For example, the importance of zfpXR in the regulation of CYP3A65 is subject to questions. It has been shown that depletion of zfpXR does not silence *cyp3a65* expression, suggesting a compensatory mechanism (Salanga et al., 2020). In contrast, two other independent studies showed that *cyp3a65* expression is not induced in zfpXR knock-out zebrafish embryos exposed to zfpXR agonist (Chang et al., 2013; Kubota et al., 2015). For zfAhR2, its role in the expression of *cyp3a65* is better defined as repression of zfAhR2 by morpholino oligonucleotides inhibits the *cyp3a65* expression and block the action of prototypical agonist of zfAhR2 such as TCDD, showing the significant role of zfAhR2 in mediating the effect of zfAhR2 agonist to induce *cyp3a65* expression (Chang et al., 2013; Tseng et al., 2005). Moreover, according to Kubota et al., zfpXR and zfAhR2 interact in the regulation of *cyp3a65* expression in a positive and reciprocal crosstalk, i.e., zfAhR2 activation promotes upregulation of *zfpXR* and *cyp3* genes, and zfpXR activation promotes upregulation of *zfahr2* and *cyp1a* genes. Such crosstalk has been suggested for human cells as AhR promotes the upregulation of *cyp3a4* and *pxr* in hepatic cell lines HepARG and HepG2, and primary hepatocytes (Kumagai et al., 2012; Rasmussen et al., 2017).

In humans, CYP3A4 is also known to be regulated by the glucocorticoid receptor (GR) in the liver and intestine (Khan et al., 2009; Usui et al., 2003). In zebrafish, there is, to our knowledge, no information regarding the role of this steroidal nuclear receptor on the regulation of *cyp3a65*, although our previous data suggest a potential role of the zfGR receptor in the developing intestine (Erradhouani et al., in submission).

In order to further characterize our transgenic model, we have undertaken the exposure of *tg(cyp3a65:GFP)* zebrafish embryos to various substances selected *a priori* from the literature data for their ability to bind and activate (agonist) or block (antagonist) the target receptor and subsequently induce or inhibit the signaling pathways mediated by the zfpXR, zfAhR2 and zfGR. For the zfpXR, agonists were clotrimazole and desogestrel and antagonist was econazole nitrate according to the results obtained *in vitro* with the HG5LN Gal4-zfpXR reporter cell line (Creusot et al., 2021). For the zfAhR2 receptor, 2,3,7,8-tetrachlorodibenzodioxin (TCDD) and 1,2,3,7,8-pentachlorodibenzo-p-dioxin (PeCDD) were selected as agonists, according to their ability to induce *cyp1a* through zfAhR2 in zebrafish embryo (Bugel et al., 2013; Goodale et al., 2012). CH223191 was selected as a zfAhR2 antagonist according to various *in vitro* mammalian models (Zhao et al., 2010), and its ability to reverse TCDD effects in zebrafish embryos (Takeda et al., 2024). In addition to these literature data, we performed an *in vitro* characterization of these agonists and antagonists in HG5LN Gal4-zfpXR and ZXLH reporter cell lines for zfpXR and zfAhR2, respectively, in order to determine whether the selected chemicals were specific for the target receptors.

For the zfGR, fluticasone propionate (FLU) was selected as a potent agonist and mifepristone (RU486) as an antagonist according to the results obtained *in vitro* with the reporter cell line UMLN-zfGR (Toso et al., 2023). Clobetasol propionate was selected as it has been characterized as a zfGR agonist *in vivo* in adult female zebrafish (Faltermann et al., 2020).

## Material & methods

### Chemicals

**Selected chemicals for zfpXR:** clotrimazole (CLO) (CAS 23,593-75-1), desogestrel (DGT) (CAS 54,024-22-5) and econazole

nitrate (ECN) (CAS 24,169-02-6). **Selected chemicals for zfAhR2:** 2,3,7,8-tetrachlorodibenzodioxin (TCDD) (CAS 1746-01-06), 1,2,3,7,8-pentachlorodibenzo-p-dioxin (PeCDD) (CAS 40,321-76-4) and CH223191 (CH22) (CAS 301,326-22-7). **Selected chemicals for zfGR:** fluticasone propionate (FLU) (CAS 80,474-14-2), clobetasol propionate (CLOBE) (CAS 25,122-46-7) and mifepristone (RU486) (CAS 84,371-65-3).

All chemicals were powders dissolved in pure dimethyl sulfoxide (DMSO) (CAS 67-68-5). All chemicals were obtained from Sigma-Aldrich (St Quentin Fallavier, France) except for TCDD, which comes from LGC Standards (Molsheim, France), and CLOBE, which comes from USP References Standard (New York City, USA).

### Animal facilities

Fish were raised in INERIS (Verneuil-en-Halatte, France) animal facilities, maintained in 3.5 L aquaria with a water recirculation system (Zebtec, Tecniplast, Italy) on a 14:10 light:dark cycle. Water quality was monitored every day: temperature ( $27.0 \pm 2.0$  °C), pH ( $7.5 \pm 0.5$ ), conductivity ( $400 \pm 50$  µS/cm), dissolved oxygen (>90%), nitrate (not detectable), nitrite (not detectable) and ammonia (not detectable). Zebrafish were fed twice a day with flake food (SDS400, Scientific Fish Food) in the morning and newly hatched brine shrimp (Sep-Art, Ocean Nutrition Europe) in the afternoon. Embryos were obtained from multiple groups of transgenic adult fish (one female and two males) placed together in specific aquaria. Eggs were laid and fertilized at night; in the morning, they were sorted and placed in an incubator at 29 °C and 14 h light/10 h dark cycle.

### Exposure of *tg(cyp3a65:GFP)* embryos

The embryos exposure concentrations were chosen from previous work in which the toxicity of the substances was tested using the fish embryo acute toxicity standardized test (OCDE, 2013). Based on these results, the concentrations of each agonist or antagonist leading to maximal GFP induction or inhibition without any apparent toxicity were selected.

Chemical exposures to zfpXR, zfAhR2 and zfGR agonists and antagonists were performed for 48 h from 72 to 120 hpf in 60 mL capacity glass crystallizers. Each condition of exposure included 3 crystallizers containing 7 embryos each. Tested substances were dissolved in breeding water at different concentrations. The control condition was DMSO 0.01% (v/v) in the case of exposure to a single substance and 0.02% (v/v) in the case of coexposure to two substances. Embryos were then exposed by balneation to the control or experimental medium with 2 mL per embryo (14 mL for 7 embryos per crystallizer) and incubated at 29 °C and 14 h light/10 h dark cycle. Water and dissolved substances were renewed every 24 h.

### *In vivo* fluorescent imaging

To quantify the GFP, living embryos were imaged by using a Zeiss AxioImager.Z1 fluorescence microscope equipped with an AxioCam Mrm camera (Zeiss GmbH, Göttingen, Germany). Embryos were placed on the slide to better capture of intestinal fluorescence. Photographs were taken with a 10 ms exposure time using a X10 objective. Data analysis was performed by Zen blue 3.3 imaging software and fluorescence quantification was carried out in the intestine by ImageJ software (Schneider et al., 2012) with a macro-control developed at the laboratory. A detection threshold of 159 ms, determined from a quantification limit in non-transgenic animals, was applied during image analysis to avoid fish autofluorescence and be specific to GFP signal.

### Quantification of mRNA expression


In addition to GFP imaging, real-time quantitative PCR (RT-qPCR)

**Table 1**

*In vitro* characterization of the tested chemicals in ZXLH and HG5LN Gal4-zfPXR cell lines. EC50 and IC50 were derived from modeled concentration-response curves using Hill regression model.

Agonists	zfAhr2 EC50 (nM)	zfPXR EC50 ( $\mu$ M)
Clotrimazole	not active	0.03
Desogestrel	not active	0.13
TCDD	0.24	not active
PeCDD	0.20	not active
Antagonists	zfAhr2 IC50 (nM)	zfPXR IC50 ( $\mu$ M)
Econazole nitrate	not active	2.8
CH223191	350	not active

was performed to assess the mRNA expressions of GFP and endogenous genes (*zfahr2*, *zfpxr*, *cyp3a65*, *cyp1a* and *fkbp5*, a zfGR target). Immediately after *in vivo* fluorescence imaging, larvae were placed in RNA stabilization solution (RNAlater™) and stored at -80 °C to preserve the samples, pending future total RNA extraction. RNA extraction, complementary DNAs generation and qPCR were performed as described in (De Oliveira et al., 2020). Primers for *zfahr2*, *zfpxr*, *cyp1a* and *fkbp5* were directly purchased from the manufacturer (ThermoFisher®), references in Supplementary Material table S1). As *cyp3a65* did not have known commercial primers, we designed them and sent the sequences to the same supplier for custom synthesis (SM table S1).



**Fig. 1.** GFP fluorescence and *cyp3a65* mRNA are induced in presence of CLO or TCDD and over-induced when embryos are coexposed to CLO+TCDD. *In vivo* imaging of tg(*cyp3a65*:GFP) embryos at 120 hpf. The dashed lines delineate the intestine. Expression in control (CTRL) DMSO 0.02% (A) and exposed embryos to CLO 1.5  $\mu$ M (B); TCDD 0.05 nM (C) and CLO 1.5  $\mu$ M + TCDD 0.05 nM (D). (E) corresponds to GFP intensity in individuals ( $n = 21$  embryos minimum per condition) after exposure to CLO 1.5  $\mu$ M (in blue); TCDD 0.05 nM (in red) and CLO 1.5  $\mu$ M + TCDD 0.05 nM (in green), control condition (CTRL) is DMSO 0.02% (in black). (F) *cyp3a65* mRNA expression in pooled zebrafish embryos ( $n = 5$  embryos/pool) in same condition.




Fig. 2. Clotrimazole-mediated zfpXR activation up-regulates zfAhr2 expression and TCDD-mediated zfAhr2 activation up-regulates zfpXR expression. zfpXR mRNA expression in pooled zebrafish embryos ( $n = 5$  embryos/pool) when exposed to (A) CLO and (C) TCDD. zfAhr2 mRNA expression in pooled zebrafish embryos ( $n = 5$  embryos/pool) when exposed to (B) CLO and (D) TCDD. Control condition (CTRL) is DMSO 0.01%.





Fig. 3. TCDD promotes strong induction of cyp1a mRNA expression when clotrimazole slightly induces it compared to TCDD. Cyp1a mRNA expression in pooled zebrafish embryos ( $n = 5$  embryos/pool) when exposed to (A) TCDD and (B) CLO.

#### *In vitro* characterization of zfpXR and zfAhr2 antagonists

***In vitro* reporter gene assay for zfAhr2** was performed in ZXLH (Zebrafish XRE Luciferase Hygromycin) cells, established from ZFL cell line (Ghosh et al., 1994) stably transfected by XRE3-TATA-Luc-Hygro construct to express the luciferase reporter gene under the transcriptional control of zfAhr2. These cells were grown at 28 °C in LDF culture medium consisting of 50% L15 medium, 35% DMEM-HG (Dulbecco's modified Eagle Medium-High Glucose), 15% Ham's F12 medium, 50

U/mL penicillin and streptomycin, 5% fetal calf serum Epidermal Growth Factor (50 ng/mL), insulin (0.01 mg/mL), HEPES (15 mM), sodium bicarbonate (0.15 g/L) and hygromycin B (0.35 μg/mL).

***In vitro* reporter gene assay for zfpXR** was performed in HG5LN Gal4-zfpXR cells, corresponding to HeLa cell line transfected using GAL4RE5-βGlobin-Luc-SVNeo construct to get HG5LN cell line, then HG5LN cell line was transfected using pSG5-GAL4(DBD)-zfpXR(LBD)-puro construct (Creusot et al., 2021). These cells were grown in Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12 (DMEM/F-12)



**Fig. 4.** GFP intensity in *tg(cyp3a65:GFP)* embryos exposed to *zfpXR* and *zfaHR2* agonist and antagonist ligands alone or in combination ( $n = 21$  embryos minimum per condition) at 120 hpf after 48 h. Exposure to CLO 1.5  $\mu\text{M}$ ; TCDD 0.05 nM; CH22 5  $\mu\text{M}$  and ECN 1  $\mu\text{M}$  alone or in combination. Control condition (CTRL) is DMSO 0.02%.

containing phenol red and 1 g/L glucose and supplemented with 5% fetal bovine serum, 100 units/mL of penicillin, 100  $\mu\text{g}/\text{mL}$  of streptomycin, 1 mg/mL geneticin and 0.5  $\mu\text{g}/\text{mL}$  puromycin in a 5%  $\text{CO}_2$  humidified atmosphere at 37  $^{\circ}\text{C}$ .

Cells were seeded in white 96-well microplates with a density of 50 000 to 100 000 cells/wells and incubated for 24 h to allow them to adhere to the plastic. Cells were then exposed to a range of concentrations of the tested molecules. Culture medium and DMSO (0.1% vol/vol) were used as negative controls. TCDD ( $10^{-8}$  mol/L) was used as a positive control for ZXLH cell line and clotrimazole ( $10^{-6}$  mol/L) for HG5LN Gal4-*zfpXR* cell line.

#### Statistics

Results were analyzed with Kruskal-Wallis and Dunn's post-hoc multiple comparisons on GraphPad Prism 8.0.2. In the figures, the number of stars corresponds to the P value summary:  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*),  $P \leq 0.001$  (\*\*\*),  $P \leq 0.0001$  (\*\*\*\*). The standard error of the arithmetic mean (SEM) is represented on each graph.

#### Results

##### *In vitro* characterization of the transactivating potential of selected agonists and antagonists

As a first step, we used well-established *in vitro* transactivation models expressing either the ligand binding domain for the *zfpXR* or the entire *zfaHR2* receptor coupled with luciferase as reporter gene to characterize the chemicals towards the two receptors investigated herein. We found that CLO selectively activated *zfpXR* with no transactivation activity on *zfaHR2*. Conversely, TCDD activated *zfaHR2* with no transactivation activity on *zfpXR* (Table 1). CH22 and ECN applied alone were unable to induce or inhibit the basal luciferase activity of both receptors, while ECN inhibited the luciferase-induced activity by CLO in *zfpXR* cell line and CH22 inhibited the TCDD-induced luciferase activity in ZXLH cell line (Table 1). These data show that the selected chemicals elicited specific activation or blocking of the target receptor and signaling pathway of interest (e.g., CLO and *zfpXR* or TCDD and *zfaHR2*) without interfering with the other receptor signaling at the cellular level (SM figure S1).

##### Crosstalk of *zfpXR* and *zfaHR2* pathways in the regulation of *cyp3a65* expression

##### Effect of each *zfpXR* or *zfaHR2* agonist alone and in combination on *cyp3a65* expression

Exposures of *tg(cyp3a65:GFP)* zebrafish embryos from 72 to 120 hpf to *zfpXR* or *zfaHR2* agonists alone promote *cyp3a65* expression as shown by induction of the GFP intensity using *in vivo* fluorescence imaging (Figs. 1B 1C and 1E) as well as induction of the relative mRNA expression using RT-qPCR (Fig. 1F). Exposure to CLO 1.5  $\mu\text{M}$  led to 3-fold induction of GFP intensity and TCDD led to 4-fold induction as compared to control (Fig. 1E). Using mRNA quantification, we show that the inductions measured in CLO or TCDD-exposed zebrafish embryos led to similar induction of the *cyp3a65* expression (Fig. 1E) as compared to GFP intensity, thereby confirming the relevance of using GFP as reported gene of the endogenous *cyp3a65* gene expression while permitting to spatialize the effect of the two compounds on the *cyp3a65* expression in the gastro-intestinal tract.







Coexposure to both CLO 1.5  $\mu\text{M}$  and TCDD 0.05 nM led to a strong induction of the *cyp3a65* expression as compared to exposure to CLO or TCDD alone as seen through *in vivo* imaging (Fig. 1D). Quantification of the GFP intensity by image analysis and *cyp3a65* expression by RT-qPCR further support this observation. The GFP intensity (11-fold induction compared to the control) and the *cyp3a65* mRNA expression (12-fold induction compared to control) in zebrafish embryos exposed to both CLO and TCDD (Figs. 1E and 1F) were also significantly different from those measured in embryos exposed to CLO or TCDD alone. To investigate if similar response profile can be observed in zebrafish embryos exposed to other agonists of *zfpXR* and *zfaHR2*, coexposure experiments were performed with DGT 1  $\mu\text{M}$  and PeCDD 0.025 nM, alone or in combination (SM figure S2). Interestingly, a similar response profile was observed, characterized by a strong induction of the GFP fluorescence and *cyp3a65* expression in the coexposure group as compared to DGT or PeCDD alone.

##### Effect of *zfpXR* and *zfaHR2* agonists on both receptors expression levels

Using the same individuals used to perform *in vivo* imaging and mRNA *cyp3a65* quantification, we also investigated the mRNA expression levels of each receptor in zebrafish embryos exposed to CLO and TCDD alone. We showed that exposure CLO, the agonist of *zfpXR* led to an up-regulation of *zfpXR* mRNA levels (Fig. 2A) but also an up-regulation of *zfaHR2* mRNA levels (Fig. 2B). In case of exposure to TCDD, the agonist of *zfaHR2*, we also measured an up-regulation of *zfaHR2* mRNA levels (Fig. 2D) as well as *zfpXR* mRNA levels (Fig. 2C).

**Table 2**

Overview of the effects of reference chemicals on *cyp3a65* expression in *tg(cyp3a65:GFP)* zebrafish embryos. Results show cooperation and ambivalent relationship between *zfpXR* and *zAhR2* signaling pathways as agonist and antagonist promote *cyp3a65* expression, and inhibition of its constitutive expression by *zfGR*.

Chemicals	Effect on <i>cyp3a65</i> expression	GFP imaging
<i>zAhR2</i> agonist	Induction +	
<i>zfpXR</i> agonist	Induction +	
<i>zAhR2</i> agonist + <i>zfpXR</i> agonist	Induction ++	
<i>zAhR2</i> agonist + <i>zfpXR</i> antagonist	Induction ++	
<i>zfpXR</i> agonist + <i>zAhR2</i> antagonist	Induction ++++	
<i>zfpXR</i> agonist	Inhibition ++++	

#### Effect of (co)exposures to *zfpXR* and *zAhR2* antagonist with CLO or TCDD

To further explore the potential interaction between *zAhR2* and *zfpXR* signaling pathways on the regulation of the *cyp3a65* gene, we then exposed zebrafish embryos to a specific and potent antagonist of *zAhR2*, *i.e.*, CH22 5  $\mu$ M, alone or in combination with CLO, the most potent agonist of *zfpXR*. Similarly, we used an antagonist of *zfpXR*, *i.e.*, ECN 1  $\mu$ M, which was tested alone or in the presence of TCDD 0.05 nM, the most potent agonist of *zAhR2*. The results showed that CH22 and ECN alone were able to induce GFP intensity as compared to control (Figs. 4A and 4B). For CH22 1  $\mu$ M, we found that at this concentration, the substance is efficient to block the action of TCDD resulting in a clear reduction of the TCDD-induced 7-ethoxyresorufin-O-deethylase activity and *cyp1a* mRNA in zebrafish embryos (SM figures S3A and S3B). Together with the data obtained in ZXLH cell line and in zebrafish

embryos, it shows that CH22 does not act as a partial agonist of *zAhR2* and is efficient in inhibiting the *zAhR2* signaling pathway. For *zfpXR*, there is, to our knowledge, no other specific *zfpXR*-regulated gene that could be investigated, and we were therefore unable to verify at the *in vivo* level the efficacy of ECN to block the *zfpXR* signaling pathway.

Coexposure to CLO and CH22 resulted in a strong induction of GFP as compared to CH22 or CLO alone (Fig. 4A). Coexposure to ECN and TCDD shows a similar trend, *i.e.*, a strong and significantly different induction of GFP as compared to ECN or TCDD alone although the magnitude of effect was less pronounced as compared to CH22 and CLO (Fig. 4B).

#### Involvement of *zfGR* pathway in the regulation of *cyp3a65* expression

Embryos exposure to *zfGR* agonist FLU promotes the mRNA expression of *fkbp5*, a well-known GR-regulated gene in zebrafish (Benato et al., 2014). This data clearly indicates that the GR-signaling pathway was activated in zebrafish exposed under this experimental condition. Interestingly, we measured a dramatic down-regulation of the *cyp3a65* mRNA expression (Figs. 5A and 5B). GFP intensity was also strongly reduced in a concentration-dependent fashion (Fig. 5C). To determine whether this reduction was restricted to FLU or can be extended to other *zfGR* agonist, we performed an exposure to another GR agonist, CLOBE, with similar results (Fig. 5D).


Considering the effects of two *zfGR* agonists on the basal expression of *cyp3a65*, we also tested *zfGR* antagonist RU486. In this experiment, TCDD was used as a positive control. As expected TCDD 0.05 nM led to a strong GFP induction, meaning that the zebrafish embryos responded in a correct manner to this treatment. RU486 alone had no effect on GFP intensity as compared to control but was significantly different as compared to FLU suggesting that *zfGR* activation is key to inhibit *cyp3a65* expression. Moreover, we observed that RU486 blocks the FLU effect on GFP intensity (Figs. 6A and 6B).

## Discussion

### New insights on the *zfpXR/zAhR2* crosstalk in *cyp3a65* expression regulation

In this study, we showed that GFP expression in *tg(cyp3a65:GFP)* is induced by different agonists of two transcription factors, *i.e.*, *zfpXR* and *zAhR2*, both being implicated in *cyp3a65* expression regulation. Combined exposure to *zfpXR* and *zAhR2* agonists promotes *cyp3a65* over-induction as compared to exposure to *zfpXR* or *zAhR2* agonists alone (Fig. 2). This indicates that both *zfpXR* and *zAhR2* signaling pathways cooperate in the regulation of *cyp3a65* expression. In addition, activation of *zfpXR* by its agonist CLO induces *zfahr2* expression, and activation of *zAhR2* by its agonist TCDD induces *zfpXR* expression (Fig. 3). It is important to remind here that we showed that TCDD is unable to act on *zfpXR* and similarly, CLO has no activity on *zAhR2* (Table 1). Conversely, at the *in vivo* level, exposure to *zfpXR* or *zAhR2* agonists led to the induction of the other signaling pathway target genes. In addition to *zfahr2* expression, we also measured *cyp1a* mRNA expression, a cytochrome known to be strongly regulated by *zAhR2* (Petrucci et al., 2001). In TCDD-exposed embryos the induction was strong. In contrast, in CLO-exposed zebrafish, the induction of *cyp1a* was weak but significant (Fig. 3). Altogether, these data are in line with Kubota et al. who showed the same crossed up-regulation between the two transcription factors in zebrafish embryos when exposed to candidate *zfpXR* agonist PCN and *zAhR2* agonist PCB126 (Kubota et al., 2015). This is also in tune with Chang et al. who assume that the interaction between the two receptors is necessary for *cyp3a65* expression (Chang et al., 2013). Such positive interaction may account for the strong induction of the *cyp3a65* gene in case of coexposure to TCDD and CLO (Fig. 1).

Surprisingly, the antagonism of *zAhR2* or *zfpXR* also promotes *cyp3a65* expression as compared to control. One hypothesis could rely on the fact that CH22 or ECN could act as partial agonists of the target



**Fig. 5.** Fluticasone propionate promotes *fkbp5* expression and inhibits *cyp3a65* expression in both GFP intensity (as clobetasol propionate) and mRNA expression. mRNA expression of (A) *fkbp5* and (B) *cyp3a65* in pooled zebrafish embryos ( $n = 5$  embryos/pool) when exposed to FLU 0.23  $\mu\text{M}$ . GFP intensity in individuals ( $n = 21$  embryos minimum per condition) after exposure to (C) FLU and (D) CLOBE. In (D) FLU is for comparison and TCDD as positive control. Control condition (CTRL) is DMSO 0.01%.


receptors. However, these two compounds were unable to activate the zfPXR or the zFAhR2 receptors *in vitro* (Table 1). Moreover, we observed a strong induction of the GFP and *cyp3a65* mRNA in case of coexposure to the antagonist of one receptor (e.g., CH22 for zFAhR2) and the agonist of the other signaling pathway (e.g., CLO for zfPXR). In such a situation, coexposure to CH22 and CLO promotes a *cyp3a65* over-induction as compared to CH22 or CLO alone (Fig. 4A). ECN and TCDD coexposure promotes *cyp3a65* over-induction as compared to ECN and TCDD alone (Fig. 4B). These results add complexity to the hypothesis of positive and reciprocal crosstalk as presented above and in literature. The interactions between the two receptors indicate that these two signaling pathways have an ambivalent relationship as they cooperate positively while in the presence of the antagonist of one receptor, the effect of the agonist of the other receptor is maximized. This ambivalent relationship could explain at least partly why in zfPXR knock-out zebrafish, *cyp3a65* appears to be induced as compared to control (Salanga et al., 2020), as in our case, the presence of CH22 or ECN led to an increased *cyp3a65* expression as compared to control. It is clear that the intimate mechanism of action between zFAhR2 and zfPXR will require additional experiments. At this stage, we can only report the complex and original interaction between zFAhR2 and zfPXR, which is highly dependent on the state of activation of each receptor to regulate the expression of *cyp3a65* in the developing intestine (Fig. 7 and Table 2).

#### New insights on zfGR involvement in *cyp3a65* expression regulation


While the zfGR agonists FLU and CLOBE promote the expression of the known zfGR-regulated gene *fkbp5*, they also inhibit *cyp3a65* expression, suggesting that zfGR-mediated pathway down-regulates the constitutive expression of *cyp3a65* (Fig. 5). This hypothesis is confirmed when embryos are exposed to both FLU and zfGR antagonist RU486; the coexposure blocks the FLU-mediated inhibition of *cyp3a65* expression (Fig. 6).

We previously performed an *in silico* analysis of the *cyp3a65* promoter which identified response elements specific to zfPXR and zFAhR2 (SM figure S4, (Erradhouani et al., in submission)), in line with the literature (Chang et al., 2013). The search for consensus motifs for zfGR did not allow us to identify specific response elements for this receptor, suggesting that zfGR modulates *cyp3a65* in another way than direct DNA binding. For example, in humans, GR can bind to the transcription factor HNF4 $\alpha$  (hepatocyte nuclear factor 4 $\alpha$ ), which binds the promoter of *cyp3a4* (Pavek et al., 2007). As GR and HNF4 $\alpha$  are well-conserved across evolution (Dinarello et al., 2020; Heppert et al., 2022); we could, therefore, pose a similar hypothesis for *cyp3a65* in zebrafish. Moreover, zfHNF4 $\alpha$  (zebrafish HNF4 $\alpha$ ) is present in the zebrafish foregut (Heppert et al., 2022), where *cyp3a65* is expressed at embryonic stage according to the *tg(cyp3a65:GFP)* model. Finally, the search for a classical direct repeat HNF4 $\alpha$  motif (DR1, 5'-AGGTCAxAGGTCA-3') led to its identification downstream of the *cyp3a65* promoter (SM figure S4). The binding





**Fig. 6.** *RU486 reverses inhibited expression of cyp3a65 by fluticasone propionate.* (A) In vivo imaging of *tg(cyp3a65:GFP)* embryos at 120 hpf after 48 h exposure to TCDD 0.05 nM; RU486 1 µM; FLU 0.23 µM and RU486+FLU. TCDD as positive control. Control condition (CTRL) is DMSO 0.02%. The dashed lines delineate the intestine. (B) GFP intensity in individuals (n = 21 embryos minimum per condition) in the same conditions.



**Fig. 7.** Hypothesis for the regulation of *cyp3a65* by zfAhR2, zfPXR and zfGR. While zfAhR2 and zfPXR link to their own response elements in the core of the *cyp3a65* promoter, the binding of zfGR to zfHNF4 $\alpha$ , itself linked to specific response elements downstream of the *cyp3a65* promoter, modulates its expression. zfAhR2 and zfPXR agonist ligands promote the expression of *cyp3a65* and zfGR inhibits its constitutive expression. XRE, PXRE and HNF-RE: response elements for zfAhR2, zfPXR and zfHNF4 $\alpha$  respectively.

of zfGR to zfHNF4 $\alpha$ , itself linked to its specific response elements (HNF-RE), could therefore modulate *cyp3a65* expression (Fig. 7). HNF4 $\alpha$  is a cofactor of PXR, allowing the expression of *cyp3a4* in humans (Tirona et al., 2003; Dvorak, 2011), it could be hypothesized that in zebrafish, the binding of zfGR to HNF4 $\alpha$  limits its role as a cofactor of zfPXR and is responsible for the observed decrease in the constitutive expression of *cyp3a65*. Overall, regulation of *cyp3a65* expression appears complex, mediated by both the ambivalent relationship existing between zfPXR and zfAhR2, and the zfGR receptor, probably through the intermediary of zfHNF4 $\alpha$  (Fig. 7 and Table 2).

If human CAR (constitutive androstane receptor) is considered “lost” in teleostean fish, human transcription factors PXR, GR and HNF4 $\alpha$  are part of the regulation of *cyp3a4* (Erradhouani et al., 2024), as zfPXR, zfGR and zfHNF4 $\alpha$  are part of the regulation of *cyp3a65*. In addition, if human AhR does not seem to be directly involved in the regulation of *cyp3a4* expression, as it is the case for zfAhR2 and *cyp3a65* in zebrafish, it could be indirectly involved by promoting PXR expression, as shown in a human hepatocyte cell line (Rasmussen et al., 2017). These similarities between human *cyp3a4* and zebrafish *cyp3a65* regulation are encouraging, in the perspective to promote our assay for MDCs assessment.

## Conclusion

In this study, we have shown that zfPXR and zfAhR2 cooperate to regulate *cyp3a65* expression. Their interactions are more complex than a simple reciprocal positive crosstalk but in contrast appear ambivalent. It is clear that results of such interactions on intestinal expression of the *cyp3a65* gene are highly dependent on the nature of the coexposure conditions.

Another major outcome of the study relies on the role of the zfGR in the constitutive expression of the *cyp3a65* gene. To our knowledge, it is the first report showing that agonists of the zfGR strongly inhibit the expression of *cyp3a65* in the developing intestine, an effect blocked in the presence of RU486, demonstrating that a functional zfGR is required in mediating the effect of zfGR agonists. We hypothesized that the mechanism through which zfGR regulates the expression of *cyp3a65* implies zfHNF4 $\alpha$  (whose responsive elements are present downstream the *cyp3a65* promoter).

By enhancing our understanding of *cyp3a65* expression regulation, we improve the *tg(cyp3a65:GFP)* model in its ability to provide information on the mode of actions of MDCs in the intestine.

## CRediT authorship contribution statement

**Chedi Erradhouani:** Writing – review & editing, Writing – original draft, Data curation, Conceptualization. **Benjamin Piccini:** Data curation. **Emmanuelle Maillot-Marechal:** Data curation. **Selim Aït-Aïssa:** Writing – review & editing. **Patrick Balaguer:** Data curation. **Xavier Coumoul:** Writing – review & editing, Conceptualization. **François Brion:** Writing – review & editing, Conceptualization.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

The authors deeply thank the anonymous reviewers for their thorough reading of the manuscript and their valuable suggestions. This study was funded by the H2020 OBERON project (grant agreement no 825712). The transgenic model was first developed within the project PestR from French National Research Program for Endocrine Disruptors (PNRPE). C.E. received a doctoral fellowship from INERIS (P190 – Ecotoxicologie). The authors acknowledge INERIS, INSERM and Université Paris-Cité for their financial and scientific support in this project, as well as the members of C.E.’s thesis committee, Dr. Etienne Blanc (Université Paris Cité) and Dr. Nicolas Diotel (Université de la Réunion).

## Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.aquatox.2025.107250](https://doi.org/10.1016/j.aquatox.2025.107250).

## Data availability

Data will be made available on request.

## References

- Audouze, K., Sarigiannis, D., Alonso-Magdalena, P., Brochot, C., Casas, M., Vrijheid, M., Babin, P.J., Karakitsios, S., Coumoul, X., Barouki, R., 2020. Integrative strategy of testing systems for identification of endocrine disruptors inducing metabolic disorders—An introduction to the OBERON project. *Int. J. Mol. Sci.* 21, 2988. <https://doi.org/10.3390/ijms21082988>.

- Benato, F., Colletti, E., Skobo, T., Moro, E., Colombo, L., Argenton, F., Dalla Valle, L., 2014. A living biosensor model to dynamically trace glucocorticoid transcriptional activity during development and adult life in zebrafish. *Mol. Cell Endocrinol.* 392, 60–72. <https://doi.org/10.1016/j.mce.2014.04.015>.
- Bernal, K., Touma, C., Erradhouani, C., Boronat-Belda, T., Gaillard, L., Al Kassir, S., Le Mentec, H., Martin-Chouly, C., Podechard, N., Lagadic-Gossmann, D., Langouet, S., Brion, F., Knoll-Gellida, A., Babin, P.J., Sovadinova, I., Babica, P., Andreau, K., Barouki, R., Vondracek, J., Alonso-Magdalena, P., Blanc, E., Kim, M.J., Coumoul, X., 2022. Combinatorial pathway disruption is a powerful approach to delineate metabolic impacts of endocrine disruptors. *FEBS Lett.* 1873–3468. <https://doi.org/10.1002/1873-3468.14465>, 14465.
- Bugel, S.M., White, L.A., Cooper, K.R., 2013. Inhibition of vitellogenin gene induction by 2,3,7,8-tetrachlorodibenzo-p-dioxin is mediated by aryl hydrocarbon receptor 2 (AHR2) in zebrafish (*Danio rerio*). *Aquatic Toxicol.* 126, 1–8. <https://doi.org/10.1016/j.aquatox.2012.09.019>.
- Chang, C.-T., Chung, H.-Y., Su, H.-T., Tseng, H.-P., Tzou, W.-S., Hu, C.-H., 2013. Regulation of zebrafish CYP3A65 transcription by AHR2. *Toxicol. Appl. Pharmacol.* 270, 174–184. <https://doi.org/10.1016/j.taap.2013.04.010>.
- Chen, J., Zhao, K.-N., Chen, C., 2014. The role of CYP3A4 in the biotransformation of bile acids and therapeutic implication for cholestasis. *Ann. Transl. Med.* 2 (7). <https://doi.org/10.3978/j.issn.2305-5839.2013.03.02>.
- Creusot, N., Garoche, C., Grimaldi, M., Boulahtouf, A., Chiavarina, B., Bourguet, W., Balaguer, P., 2021. A comparative study of human and zebrafish pregnane X receptor activities of pesticides and steroids using in vitro reporter gene assays. *Front. Endocrinol.* 12, 665521. <https://doi.org/10.3389/fendo.2021.665521>.
- De Oliveira, J., Chadili, E., Piccini, B., Turies, C., Maillot-Maréchal, E., Palluel, O., Pardon, P., Budzinski, H., Cousin, X., Brion, F., Hinfray, N., 2020. Refinement of an OECD test guideline for evaluating the effects of endocrine disrupting chemicals on aromatase gene expression and reproduction using novel transgenic cyp19a1a-eGFP zebrafish. *Aquatic Toxicol.* 220, 105403. <https://doi.org/10.1016/j.aquatox.2020.105403>.
- Dinarello, A., Licciardello, G., Fontana, C.M., Tiso, N., Argenton, F., Dalla Valle, L., 2020. Glucocorticoid receptor activities in the zebrafish model: a review. *J. Endocrinol.* 247, R63–R82. <https://doi.org/10.1530/JOE-20-0173>.
- Dvorak, Z., 2011. Drug–drug interactions by azole antifungals: beyond a dogma of CYP3A4 enzyme activity inhibition. *Toxicol. Lett.* 202, 129–132. <https://doi.org/10.1016/j.toxlet.2011.01.027>.
- Erradhouani, C., Bortoli, S., Ait-Aïssa, S., Coumoul, X., Brion, F., 2024. Metabolic disrupting chemicals in the intestine: the need for biologically relevant models. *FEBS. Open. Bio* 14, 1397–1419. <https://doi.org/10.1002/2211-5463.13878>.
- Erradhouani, C., Geffroy, F., Piccini, B., Hinfray, N., Sohm, F., Chadili, E., Balaguer, P., Ait-Aïssa, S., Coumoul, X., Brion, F., in submission. EMERGE, a new zebrafish embryo *in vivo* assay to assess metabolic endocrine-disrupting chemicals in the intestine. In submission.
- Faltermann, S., Hettich, T., Küng, N., Fent, K., 2020. Effects of the glucocorticoid clobetasol propionate and its mixture with cortisol and different class steroids in adult female zebrafish. *Aquatic Toxicol.* 218, 105372. <https://doi.org/10.1016/j.aquatox.2019.105372>.
- Ghosh, C., Zhou, Y.L., Collodi, P., 1994. Derivation and characterization of a zebrafish liver cell line. *Cell Biol. Toxicol.* 10, 167–176. <https://doi.org/10.1007/BF00757560>.
- Goldstone, J.V., McArthur, A.G., Kubota, A., Zanette, J., Parente, T., Jönsson, M.E., Nelson, D.R., Stegeman, J.J., 2010. Identification and developmental expression of the full complement of Cytochrome P450 genes in Zebrafish. *BMC. Genomics.* 11, 643. <https://doi.org/10.1186/1471-2164-11-643>.
- Goodale, B.C., La Du, J.K., Bisson, W.H., Janszen, D.B., Waters, K.M., Tanguay, R.L., 2012. AHR2 mutant reveals functional diversity of aryl hydrocarbon receptors in Zebrafish. *PLoS. One* 7, e29346. <https://doi.org/10.1371/journal.pone.0029346>.
- Heppert, J.K., Lickwar, C.R., Tillman, M.C., Davis, B.R., Davison, J.M., Lu, H.-Y., Chen, W., Busch-Nentwich, E.M., Corcoran, D.L., Rawls, J.F., 2022. Conserved roles for Hnf4 family transcription factors in zebrafish development and intestinal function. *Genetics* 222, iyac133. <https://doi.org/10.1093/genetics/iyac133>.
- Kassotis, C.D., Vom Saal, F.S., Babin, P.J., Lagadic-Gossmann, D., Le Mentec, H., Blumberg, B., Mohajer, N., Legrand, A., Munic Kos, V., Martin-Chouly, C., Podechard, N., Langouët, S., Touma, C., Barouki, R., Kim, M.J., Audouze, K., Choudhury, M., Shree, N., Bansal, A., Howard, S., Heindel, J.J., 2022. Obesity III: obesogen assays: limitations, strengths, and new directions. *Biochem. Pharmacol.* 199, 115014. <https://doi.org/10.1016/j.bcp.2022.115014>.
- Khan, A.A., Chow, E.C.Y., Van Loenen-Weemaes, A.M.A., Porte, R.J., Pang, K.S., Groothuis, G.M.M., 2009. Comparison of effects of VDR versus PXR, FXR and GR ligands on the regulation of CYP3A isozymes in rat and human intestine and liver. *Europ. J. Pharmaceut. Sci.* 37, 115–125. <https://doi.org/10.1016/j.ejps.2009.01.006>.
- Kubota, A., Goldstone, J.V., Lemaire, B., Takata, M., Woodin, B.R., Stegeman, J.J., 2015. Role of Pregnane X receptor and Aryl hydrocarbon receptor in transcriptional regulation of pxxr, CYP2, and CYP3 genes in developing zebrafish. *Toxicol. Sci.* 143, 398–407. <https://doi.org/10.1093/toxsci/kfu240>.
- Kumagai, T., Suzuki, H., Sasaki, T., Sakaguchi, S., Miyairi, S., Yamazoe, Y., Nagata, K., 2012. Polycyclic aromatic hydrocarbons activate CYP3A4 gene transcription through human pregnane X receptor. *Drug Metab. Pharmacokinet.* 27, 200–206. <https://doi.org/10.2133/dmpk.DMPK-11-RG-094>.
- OCDE, 2013. Test No. 236: fish embryo acute toxicity (FET) Test, OECD guidelines for the testing of chemicals, section 2. OECD. <https://doi.org/10.1787/9789264203709-en>.
- Pavek, P., Cerveny, L., Svecova, L., Brysch, M., Libra, A., Vrzal, R., Nachtigal, P., Staud, F., Ulrichova, J., Fendrich, Z., Dvorak, Z., 2007. Examination of Glucocorticoid receptor alpha-mediated transcriptional regulation of P-glycoprotein, CYP3A4, and CYP2C9 genes in placental trophoblast cell lines. *Placenta* 28, 1004–1011. <https://doi.org/10.1016/j.placenta.2007.05.001>.
- Petrakis, D., Vassilopoulou, L., Mamoulakis, C., Psycharakis, C., Anifantaki, A., Sifakis, S., Docea, A., Tsiaoussis, J., Makriannakis, A., Tsatsakis, A., 2017. Endocrine disruptors leading to obesity and related diseases. *IJERPH* 14, 1282. <https://doi.org/10.3390/ijerph14101282>.
- Petrulis, J.R., Chen, G., Bann, S., LaMarre, J., Bunce, N.J., 2001. Application of the ethoxoresorufin-O-deethylase (EROD) assay to mixtures of halogenated aromatic compounds. *Environ. Toxicol.* 16, 177–184. <https://doi.org/10.1002/tox.1022>.
- Rasmussen, M.K., Daujat-Chavanieu, M., Gerbal-Chaloin, S., 2017. Activation of the aryl hydrocarbon receptor decreases rifampicin-induced CYP3A4 expression in primary human hepatocytes and HepaRG. *Toxicol. Lett.* 277, 1–8. <https://doi.org/10.1016/j.toxlet.2017.05.029>.
- Salanga, M.C., Brun, N.R., Francolini, R.D., Stegeman, J.J., Goldstone, J.V., 2020. CRISPR-Cas9-Mutated Pregnane X Receptor (pxr) Retains Pregnenolone-induced Expression of cyp3a65 in Zebrafish (*Danio rerio*) Larvae. *Toxicol. Sci.* 174, 51–62. <https://doi.org/10.1093/toxsci/kfz246>.
- Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* 9, 671–675. <https://doi.org/10.1038/nmeth.2089>.
- Takeda, K., Sarata, A., Terasaki, M., Kubota, A., Shimizu, K., Kamata, R., 2024. Assessment of the aryl hydrocarbon receptor-mediated effects of aromatic sensitizers in paper recycling effluent employing zebrafish embryos and in silico docking. *Environ. Toxicol. Chem.* <https://doi.org/10.1002/etc.5969>.
- Tirona, R.G., Lee, W., Leake, B.F., Lan, L.-B., Cline, C.B., Lamba, V., Parviz, F., Duncan, S. A., Inoue, Y., Gonzalez, F.J., Schuetz, E.G., Kim, R.B., 2003. The orphan nuclear receptor HNF4 $\alpha$  determines PXR- and CAR-mediated xenobiotic induction of CYP3A4. *Nat. Med.* 9, 220–224. <https://doi.org/10.1038/nm815>.
- Toso, A., Boulahtouf, A., Escande, A., Garoche, C., Balaguer, P., 2023. A comparative study of human and zebrafish glucocorticoid receptor activities of natural and pharmaceutical steroids. *Front. Endocrinol.* 14, 1235501. <https://doi.org/10.3389/fendo.2023.1235501>.
- Tseng, H., Hseu, T., Buhler, D., Wang, W., Hu, C., 2005. Constitutive and xenobiotics-induced expression of a novel CYP3A gene from zebrafish larva. *Toxicol. Appl. Pharmacol.* 205, 247–258. <https://doi.org/10.1016/j.taap.2004.10.019>.
- Usui, T., Saitoh, Y., Komada, F., 2003. Induction of CYP3As in HepG2 cells by several drugs.—Association between induction of CYP3A4 and expression of glucocorticoid receptor—26.
- van Herwaarden, A.E., van Waterschoot, R.A.B., Schinkel, A.H., 2009. How important is intestinal cytochrome P450 3A metabolism? *Trends. Pharmacol. Sci.* 30, 223–227. <https://doi.org/10.1016/j.tips.2009.02.003>.
- van Waterschoot, R.A.B., Rooswinkel, R.W., Wagenaar, E., van der Kruijssen, C.M.M., van Herwaarden, A.E., Schinkel, A.H., 2009. Intestinal cytochrome P450 3A plays an important role in the regulation of detoxifying systems in the liver. *The FASEB J.* 23, 224–231. <https://doi.org/10.1096/fj.08-114876>.
- White, R.M., Sessa, A., Burke, C., Bowman, T., LeBlanc, J., Ceol, C., Bourque, C., Dovey, M., Goessling, W., Burns, C.E., Zon, L.I., 2008. Transparent adult zebrafish as a tool for in vivo transplantation analysis. *Cell Stem Cell* 2, 183–189. <https://doi.org/10.1016/j.stem.2007.11.002>.
- Zhao, B., DeGroot, D.E., Hayashi, A., He, G., Denison, M.S., 2010. CH223191 is a ligand-selective antagonist of the Ah (Dioxin) receptor. *Toxicol. Sci.* 117, 393–403. <https://doi.org/10.1093/toxsci/kfq217>.